

Sjögren Syndrome

Sjögren syndrome is an **autoimmune disorder** in which the body's white blood cells attack and destroy other cells. It can exist by itself (primary Sjögren syndrome) or develop in association with another disorder such as rheumatoid arthritis, systemic sclerosis, primary biliary cirrhosis, or Hashimoto thyroiditis (associated Sjögren syndrome). Hallmarks are the dry mouth and dry eyes known as the **sicca syndrome**. Sjögren syndrome affects 1 million to 4 million people in the United States. Most are over 40 years old at the time of diagnosis. Women are 9 times more likely to have Sjögren syndrome than men. The disease is named after Swedish **ophthalmologist** (eye physician) Henrik Sjögren (1899-1986), who first described it. The July 28, 2010, issue of *JAMA* includes an article about treatment of primary Sjögren syndrome.

SYMPTOMS

Although Sjögren syndrome often presents only with the sicca syndrome, symptoms can occur in any part of the body. Dryness of the eyes may severely damage the cornea, resulting in a scratchy or irritated sensation. Lack of tears can cause permanent eye damage. Insufficient saliva can cause cavities and dull taste and smell and make eating and swallowing painful. Dryness of the trachea can result in cough. In one-third of people with Sjögren syndrome, the **parotid** (salivary) glands become enlarged and tender. One-third of patients may develop joint pain. Lymph nodes throughout the body may enlarge. Nerves, lungs, and other organs may be damaged by inflammation. Debilitating fatigue and pain can impair quality of life.

DIAGNOSIS

A sensation of dry mouth and dry eyes with joint inflammation is often sufficient to suspect the diagnosis. Tear production is tested for by the **Schirmer test** (a filter paper strip is placed inside each lower eyelid for 5 minutes). Ophthalmologists test for damage to the **cornea** (eye surface) using fluorescein staining. A salivary gland biopsy may be required to confirm the diagnosis. Your doctor may also perform some blood tests or other studies.


TREATMENTS

There is no specific treatment for Sjögren syndrome. Treatment of sicca manifestations is mainly symptomatic with saliva substitutes and eyedrops and is typically intended to limit the damage that can result from chronic dryness. Nonsteroidal anti-inflammatory drugs such as ibuprofen may be used to treat musculoskeletal symptoms. People with severe symptoms are treated with steroids or immunosuppressive drugs. New medications such as monoclonal antibodies are under investigation.


PROGNOSIS

Prognosis in Sjögren syndrome varies. Many patients have mild disease, experiencing only dry eyes and mouth; others go through cycles of good health followed by severe disease. At any time, disease activity can remain stable, worsen, or go into remission. People with Sjögren syndrome have a higher rate of B-cell lymphoma; those with severe disease are more likely to develop lymphoma. Overall, 5% of people with Sjögren syndrome develop some form of malignancy.

Tests for Sjögren Syndrome
Schirmer test



Fluorescein corneal staining
(performed in dark room)



FOR MORE INFORMATION

- National Institute of Neurological Disorders and Stroke
www.ninds.nih.gov/disorders/sjogrens
- Sjögren Syndrome Foundation
www.sjogrens.org

INFORM YOURSELF

To find this and previous JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. Many are available in English and Spanish.

Sources: National Institute of Neurological Disorders and Stroke, Sjögren Syndrome Foundation

Huan J. Chang, MD, MPH, Writer

Alison E. Burke, MA, Illustrator

Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 312/464-0776.

